

Exclusive Semileptonic Rare Decays $B \rightarrow K^{(*)}l^+l^-$ in a SUSY SO(10) GUT

Wen-Jun Li^{a,b}, Yuan-Ben Dai^a, Chao-Shang Huang^a

^a*Institute of Theoretical Physics, Academia Sinica, P. O. Box 2735, Beijing 100080, China*

^b*Graduate School of the Chinese Academy of Science, YuQuan Road 19A, Beijing 100039, China*

In the SUSY SO(10) GUT context, we study the exclusive processes $B \rightarrow K^{(*)}l^+l^-$ ($l = \mu, \tau$). Using the Wilson coefficients of relevant operators including the new operators $Q'_{1,2}$ which are induced by neutral Higgs boson (NHB) penguins, we evaluate some possible observables associated with these processes like, the invariant mass spectrum (IMS), lepton pair forward backward asymmetry (FBA), lepton polarization asymmetries etc. In the model the contributions from Wilson coefficients $C'_{Q_{1,2}}$, among new contributions, are dominant. Our results show that the NHB effects are sensitive to the FBA, $dL/d\hat{s}$, and $dT/d\hat{s}$ of $B \rightarrow K^{(*)}\tau^+\tau^-$ decay, which are expected to be measured in B factories, the deviation of $dT/d\hat{s}$ in $B \rightarrow K\mu^+\mu^-$ can reach 0.1 from SM, which could be seen in B factories, and the average of the normal polarization $dN/d\hat{s}$ can reach several percent for $B \rightarrow K\mu^+\mu^-$ and it is 0.05 or so for $B \rightarrow K\tau^+\tau^-$, which could be measured in the future super B factories and provide a useful information to probe new physics and discriminate different models.

I. INTRODUCTION

The rich flavor changing neutral current processes $B \rightarrow K^{(*)}l^+l^-$ have been the sharper focus since these decays are potential testing grounds for the SM at loop level and hoped to probe the new physics beyond the SM. Recently exclusive measurements have been done by Belle and BaBar and the following results for the branching ratios of the $B \rightarrow K\ell^+\ell^-$ and $B \rightarrow K^*\ell^+\ell^-$ ($\ell = e, \mu$) decays are announced [1, 2]:

$$\begin{aligned} Br(B \rightarrow K\ell^+\ell^-) &= \begin{cases} (4.8^{+1.0}_{-0.9} \pm 0.3 \pm 0.1) \times 10^{-7} & [Belle], \\ (0.65^{+0.14}_{-0.13} \pm 0.04) \times 10^{-6} & [BaBar], \end{cases} \\ Br(B \rightarrow K^*\ell^+\ell^-) &= \begin{cases} (11.5^{+2.6}_{-2.4} \pm 0.8 \pm 0.2) \times 10^{-7} & [Belle], \\ (0.88^{+0.33}_{-0.29}) \times 10^{-6} & [BaBar], \end{cases} \end{aligned} \quad (1)$$

which imply

$$\begin{aligned} Br(B \rightarrow K\ell^+\ell^-)_{\text{world-ave}} &= (0.54 \pm 0.09) \times 10^{-6} \\ Br(B \rightarrow K^*\ell^+\ell^-)_{\text{world-ave}} &= (1.04 \pm 0.22) \times 10^{-6} \end{aligned} \quad (2)$$

In addition, the FBA for $B \rightarrow K^*\ell^+\ell^-$ have firstly observed at Belle collider [1].

The $B \rightarrow K^{(*)}l^+l^-$ decays, induced by $b \rightarrow sl^+l^-$ transition at quark level, are experimentally easier to measure than inclusive processes $B \rightarrow X_sl^+l^-$. From the theoretical point of view there are large uncertainties, which come mostly from the decay form factors, to make predictions for the exclusive processes. At present, the knowledge of form factors lacks a precisely non-perturbative solution. A number of papers are dedicated to calculating the form factors with various appropriate methods [3, 4, 5, 6]. Among them, the QCD sum rules on the light-cone (LCSRs), which deals with form factors at small values of $\hat{s}$, the momentum transfer to leptons, is a complementary to lattice approach and has the consistence with perturbative QCD and the heavy quark limit. In this paper, we will use the form factors calculated by the LCSRs [14].

The measurement of invariant mass spectrum, forward-backward asymmetry, and lepton polarizations are efficient tools to establish the new physics. There are a great deal of studies for processes $B \rightarrow K^{(*)}l^+l^-$ in theory. A model independent analysis has been carried out in [7, 8] and a lot of papers perform the investigation in many new physics scenarios [9, 10, 11, 12, 13, 14, 15, 16, 17, 18] and some works [19, 20, 21] are dedicated to the double lepton polarization. It has been pointed out [12, 13, 16] that in the some types of two-Higgs-double model and SUSY models the neutral Higgs bosons have sizable contributions to these decays (for $l = \mu, \tau$) at large $\tan\beta$. In [14], Ali *et.al* calculate these quantities in five scenarios of supersymmetric model with assuming no additional phase. Kruger's studies [15] are focus on the CP violation of $\bar{B} \rightarrow K\ell^+l^-$ in the model with additional CP phases and an extended operator basis. In ref. [13] only the Higgs penguins with chargino-stop propagated in the loop are considered.

The rapid progress in neutrino experiment [22] requires new physics to provide a theoretical explanation. Motivated by neutrino observations, a number of SUSY SO(10) models have been proposed [23, 24, 25, 26] and some phenomenological consequences of the models have been discussed [23, 27, 28, 29, 30]. In SUSY SO(10) GUT models,

there is a complex flavor non-diagonal down-type squark mass matrix element of 2nd and 3rd generations of order one in the RR sector (i.e., δ_{23}^{dRR} non-zero with $\delta_{23}^{dRR} \equiv (M_{dRR}^2)_{23}/m_{\tilde{q}}^2$, where $(M_{dRR}^2)_{ij}$ is the flavor non-diagonal squared right-handed down squark mass matrix element and $m_{\tilde{q}}$ is the average right-handed down-type squark mass) at the GUT scale [25] which can induce large flavor off-diagonal couplings such as the coupling of gluino to the quark and squark which belong to different generations. These couplings are in general complex and may contribute to the process of flavor changing neutral currents (FCNC). For specific, we use the SUSY SO(10) model described in ref. [25]. The details and a simple description of this model can be found in Refs. [25, 30]. In this paper, we investigate exclusive decay $B \rightarrow K^{(*)}l^+l^-$ ($l = \mu, \tau$) in the context of SUSY SO(10) GUT. It is well-known that the effects of the counterparts of usual chromo-magnetic and electro-magnetic dipole moment operators as well as semileptonic operators with opposite chirality are suppressed by m_s/m_b and consequently negligible in SM. However, in SUSY SO(10) GUTs their effects can be significant, since δ_{23}^{dRR} can be as large as 0.5 [25]. Furthermore, δ_{23}^{dRR} can induce new operators, the counterparts of usual scalar operators ($Q_{1,2}$, for their definitions, see below) in SUSY models, due to NHB penguins with gluino-down type squark propagated in the loop. We include the contributions of these counterpart operators and find that indeed they are dominant in the SUSY SO(10), using the MIA with double insertions to calculate Wilson coefficients of operators. The aim of our paper is make an analysis of the SUSY contributions, in particular, the contributions of neutral Higgs bosons, to the exclusive decay $B \rightarrow K^{(*)}l^+l^-$ ($l = \mu, \tau$) in the context of SUSY SO(10) GUT.

The paper is organized as follows. In section 2, we present the effective Hamiltonian and hadronic matrix elements of relevant operators in terms of form factors. In section 3, the expressions of observables are given. In section 4, we give the sparticle mass spectrum using the revised ISAJET. We make the numerical analysis and draw the conclusion in section 5.

II. EFFECTIVE HAMILTONIAN AND FORM FACTORS

In the SUSY SO(10) GUT, after integrating the heavy degree of freedom from the full theory, the general effective Hamiltonian for $b \rightarrow sl^+l^-$ can be written as follows:

$$\mathcal{H}_{\text{eff}} = -\frac{4G_F}{\sqrt{2}}V_{tb}V_{ts}^* \left[\sum_{i=1}^2 C_i(\mu) O_i(\mu) + \sum_{i=3}^{10} (C_i(\mu) O_i(\mu) + C'_i(\mu) O'_i(\mu)) + \sum_{i=1}^8 (C_i(\mu) Q_i(\mu) + C'_i(\mu) Q'_i(\mu)) \right] \quad (3)$$

where $O_i(\mu)$ ($i = 1, \dots, 10$) are dimension-six operators and $C_i(\mu)$ are the corresponding Wilson coefficients at the scale μ [31]. The additional operators Q_i ($i = 1, \dots, 8$) come from the neutral Higgs exchange diagrams and their definitions are given as [16, 32]

$$\begin{aligned} Q_1 &= \frac{e^2}{16\pi^2} (\bar{s}_L^\alpha b_R^\alpha) (\bar{l}l) \\ Q_2 &= \frac{e^2}{16\pi^2} (\bar{s}_L^\alpha b_R^\alpha) (\bar{l}\gamma_5 l) \\ Q_{3(4)} &= \frac{g^2}{16\pi^2} (\bar{s}_L^\alpha b_R^\alpha) \left(\sum_q \bar{q}_{L(R)}^\beta q_{R(L)}^\beta \right) \\ Q_{5(6)} &= \frac{g^2}{16\pi^2} (\bar{s}_L^\alpha b_R^\beta) \left(\sum_q \bar{q}_{L(R)}^\beta q_{R(L)}^\alpha \right) \\ Q_7 &= \frac{g^2}{16\pi^2} (\bar{s}_L^\alpha \sigma^{\mu\nu} b_R^\alpha) \left(\sum_q \bar{q}_L^\beta \sigma_{\mu\nu} q_R^\beta \right) \\ Q_8 &= \frac{g^2}{16\pi^2} (\bar{s}_L^\alpha \sigma^{\mu\nu} b_R^\beta) \left(\sum_q \bar{q}_L^\beta \sigma_{\mu\nu} q_R^\alpha \right) \end{aligned} \quad (4)$$

and the corresponding Wilson coefficients can be found in [33]. The primed operators, the counterpart of the unprimed operators, are obtained by replacing the chiralities in the corresponding unprimed operators with opposite ones. The explicit expressions of the operators governing $B \rightarrow K^{(*)}l^+l^-$ are given as:

$$\begin{aligned} O_7 &= \frac{e}{16\pi^2} m_b (\bar{s} \sigma_{\mu\nu} P_R b) F^{\mu\nu}, & O'_7 &= \frac{e}{16\pi^2} m_b (\bar{s} \sigma_{\mu\nu} P_L b) F^{\mu\nu}, \\ O_9 &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_L b) (\bar{l} \gamma^\mu l), & O'_9 &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_R b) (\bar{l} \gamma^\mu l), \end{aligned}$$

$$\begin{aligned}
O_{10} &= \frac{e^2}{16\pi^2}(\bar{s}\gamma_\mu P_L b)(\bar{l}\gamma^\mu \gamma_5 l), & O'_{10} &= \frac{e^2}{16\pi^2}(\bar{s}\gamma_\mu P_R b)(\bar{l}\gamma^\mu \gamma_5 l), \\
Q_1 &= \frac{e^2}{16\pi^2}(\bar{s}P_R b)(\bar{l}l), & Q'_1 &= \frac{e^2}{16\pi^2}(\bar{s}P_L b)(\bar{l}l), \\
Q_2 &= \frac{e^2}{16\pi^2}(\bar{s}P_R b)(\bar{l}\gamma_5 l), & Q'_2 &= \frac{e^2}{16\pi^2}(\bar{s}P_L b)(\bar{l}\gamma_5 l)
\end{aligned} \tag{5}$$

where $P_{L,R} = (1 \mp \gamma_5)/2$. From the above Hamiltonian, we get the decay amplitude of $b \rightarrow sl^+l^-$:

$$\begin{aligned}
\mathcal{M}(b \rightarrow sl^+l^-) = & -\frac{G_F\alpha}{\sqrt{2}\pi}V_{tb}V_{ts}^* \left\{ C_9^{eff}[\bar{s}\gamma_\mu Lb][\bar{l}\gamma^\mu l] + C_{10}[\bar{s}\gamma_\mu Lb][\bar{l}\gamma^\mu \gamma_5 l] \right. \\
& \left. - 2\hat{m}_b C_7^{eff}[\bar{s}i\sigma_{\mu\nu}\frac{\hat{q}^\nu}{\hat{s}}Rb][\bar{l}\gamma^\mu l] + C_{Q_1}[\bar{s}Rb][\bar{l}l] + C_{Q_2}[\bar{s}Rb][\bar{l}\gamma_5 l] + (C_i(m_b) \leftrightarrow C'_i(m_b)) \right\} \tag{6}
\end{aligned}$$

where $s = q^2$, $\hat{s} = \frac{s}{m_B^2}$, $q = p_B - p_{K(*)}$ is the momentum transfer. The Wilson coefficient $C_9^{eff}(\mu)$ and C_7^{eff} are defined as:

$$C_9^{eff}(\mu, \hat{s}) = C_9(\mu) + Y(\mu, \hat{s}) + \frac{3\pi}{\alpha^2}C(\mu)\Sigma_{V_i=\psi(1s)\dots\psi(6s)}^{k_i}\frac{\Gamma(V_i \rightarrow l^+l^-)m_{V_i}}{m_{V_i}^2 - \hat{s}m_B^2 - im_{V_i}\Gamma_{V_i}}, \tag{7}$$

$$C_7^{eff} = C_7 - C_5/3 - C_6 \tag{8}$$

where $C_9^{eff}(\mu)$ contains the long distance effects associated with real $\bar{c}c$ in the intermediate states $B \rightarrow KJ/\psi(\psi') \rightarrow Kl^+l^-$, which can be expressed as the last term in Eq.(7), as well as the short distance contributions. The function $Y(\mu, \hat{s})$ comes from the one-loop contributions of the four-quark operators and its explicit expression can be found in [34]. The $C_9^{eff}(\mu)$ and C_7^{eff} can be obtained by replacing the unprimed Wilson coefficients with the corresponding primed ones in the above formula.

In virtue of the form factors in [14], the hadronic matrix elements in the $B \rightarrow Kl^+l^-$ decay can be expressed as:

$$\langle K(p)|\bar{s}\gamma_\mu b|B(p_b) \rangle = f_+(s)p_\mu + f_-(s)q_\mu, \tag{9}$$

$$\langle K(p)|\bar{s}\sigma_{\mu\nu}q^\nu(1 + \gamma_5)b|B(p_b) \rangle = i\{p_\mu s - q_\mu(m_B^2 - m_K^2)\}\frac{f_T(s)}{m_B + m_K} \tag{10}$$

Using equations of motion, we obtain

$$\langle K(p_K)|\bar{s}b|B(p_B) \rangle = \frac{m_B^2 - m_K^2}{m_s - m_b}f_0(s). \tag{11}$$

For $B \rightarrow K^*l^+l^-$, the form factors are defined as follows.

$$\begin{aligned}
\langle K^*(p_{K^*}, \varepsilon)|\bar{s}\gamma_\mu(1 \pm \gamma_5)b|B(p_B) \rangle &= \epsilon_{\mu\nu\rho\sigma}\epsilon^{*\nu}p_B^\rho p_{K^*}^\sigma \frac{2V(s)}{m_B + m_{K^*}} \pm i\epsilon_\mu^*(m_B + m_{K^*})A_1(s) \\
&\mp ip_\mu(\epsilon^*p_B)\frac{A_2(s)}{m_B + m_{K^*}} \\
&\mp iq_\mu(\epsilon^*p_B)\frac{2m_{K^*}}{s}(A_3(s) - A_0(s)),
\end{aligned} \tag{12}$$

$$\begin{aligned}
\langle K^*(p_{K^*}, \varepsilon)|\bar{s}\sigma_{\mu\nu}q^\nu(1 \pm \gamma_5)b|B(p_B) \rangle &= i\epsilon_{\mu\nu\rho\sigma}\epsilon^{*\nu}p_B^\rho p_{K^*}^\sigma 2T_1(s) \pm \epsilon_\mu^*T_2(s)(m_B^2 - m_{K^*}^2) \\
&\mp (\epsilon^*p_B)p_\mu \left(T_2(s) + T_3(s)\frac{s}{m_B^2 - m_{K^*}^2} \right) \pm (\epsilon^*p_B)q_\mu T_3(s)
\end{aligned} \tag{13}$$

and

$$\langle K^*(p_{K^*}, \varepsilon)|\bar{s}(1 \pm \gamma_5)b|B(p_B) \rangle = \mp i(\epsilon^*p_B)\frac{2m_{K^*}}{m_b + m_s}A_0(s) \tag{14}$$

by means of equations of motion.

The q^2 dependence of the form factors can be parameterized as

$$F(\hat{s}) = F(0)\exp(c_1\hat{s} + c_2\hat{s}^2 + c_3\hat{s}^3)$$

where related parameters are given in the Table.4 of [14].

III. THE FORMULA FOR OBSERVABLES

From Eq.(3-14), we can write the decay matrix elements as

$$\mathcal{A} = -\frac{G_F \alpha}{2\sqrt{2}\pi} V_{tb} V_{ts}^* m_B [T_\mu^1 (\bar{l} \gamma^\mu l) + T_\mu^2 (\bar{l} \gamma^\mu \gamma_5 l) + S(\bar{l} l)] \quad (15)$$

where for $B \rightarrow Kl^+l^-$ decay:

$$\begin{aligned} T_\mu^1 &= A'(\hat{s}) \hat{p}_\mu, \\ T_\mu^2 &= C'(\hat{s}) \hat{p}_\mu + D'(\hat{s}) \hat{q}_\mu, \\ S &= S_1(\hat{s}) \end{aligned} \quad (16)$$

and for $B \rightarrow K^*l^+l^-$ decay:

$$\begin{aligned} T_\mu^1 &= A(\hat{s}) \epsilon_{\mu\rho\alpha\beta} \epsilon^{*\rho} \hat{p}_B^\alpha \hat{p}_{K^*}^\beta - iB(\hat{s}) \epsilon_\mu^* + iC(\hat{s}) (\epsilon^* \cdot \hat{p}_B) \hat{p}_\mu, \\ T_\mu^2 &= E(\hat{s}) \epsilon_{\mu\rho\alpha\beta} \epsilon^{*\rho} \hat{p}_B^\alpha \hat{p}_{K^*}^\beta - iF(\hat{s}) \epsilon_\mu^* + iG(\hat{s}) (\epsilon^* \cdot \hat{p}_B) \hat{p}_\mu + iH(\hat{s}) (\epsilon^* \cdot \hat{p}_B) \hat{q}_\mu, \\ S &= i2\hat{m}_{K^*} (\epsilon^* \cdot \hat{p}_B) S_2(\hat{s}) \end{aligned} \quad (17)$$

where $p = p_B + p_{K^*}$, $q = p_B - p_{K^*}$, $\hat{m} = \frac{m}{m_B}$, $\hat{p} = \frac{p}{m_B}$, and the auxiliary functions are defined as:

$$A'(\hat{s}) = [C_9^{eff}(\hat{s}) + C_9'^{eff}(\hat{s})] f_+(\hat{s}) + \frac{2\hat{m}_b}{1 + \hat{m}_K} (C_7^{eff} + C_7'^{eff}) f_T(\hat{s}), \quad (18)$$

$$C'(\hat{s}) = (C_{10} + C_{10}') f_+(\hat{s}), \quad (19)$$

$$D'(\hat{s}) = (C_{10} + C_{10}') f_-(\hat{s}) - \frac{1 - \hat{m}_K^2}{2\hat{m}_l(\hat{m}_b - \hat{m}_s)} (C_{Q_2} + C_{Q_2}') f_0(\hat{s}), \quad (20)$$

$$S_1(\hat{s}) = \frac{1 - \hat{m}_K^2}{(\hat{m}_b - \hat{m}_s)} (C_{Q_1} + C_{Q_1}') f_0(\hat{s}), \quad (21)$$

$$A(\hat{s}) = \frac{2V(\hat{s})}{1 + \hat{m}_{K^*}} [C_9^{eff}(\hat{s}) + C_9'^{eff}(\hat{s})] + \frac{4\hat{m}_b}{\hat{s}} (C_7^{eff} + C_7'^{eff}) T_1(\hat{s}), \quad (22)$$

$$B(\hat{s}) = (1 + \hat{m}_{K^*}) [C_9^{eff}(\hat{s}) - C_9'^{eff}(\hat{s})] A_1(\hat{s}) + \frac{2\hat{m}_b}{\hat{s}} (1 - \hat{m}_{K^*}^2) (C_7^{eff} - C_7'^{eff}) T_2(\hat{s}), \quad (23)$$

$$C(\hat{s}) = \frac{A_2(\hat{s})}{(1 + \hat{m}_{K^*})} [C_9^{eff}(\hat{s}) - C_9'^{eff}(\hat{s})] + \frac{2\hat{m}_b}{1 - \hat{m}_{K^*}^2} (C_7^{eff} - C_7'^{eff}) \left(T_3(\hat{s}) + \frac{1 - \hat{m}_{K^*}^2}{\hat{s}} T_2(\hat{s}) \right), \quad (24)$$

$$E(\hat{s}) = \frac{2V(\hat{s})}{1 + \hat{m}_{K^*}} (C_{10} + C_{10}'), \quad (25)$$

$$F(\hat{s}) = (1 + \hat{m}_{K^*}) (C_{10} - C_{10}') A_1(\hat{s}), \quad (26)$$

$$G(\hat{s}) = \frac{1}{1 + \hat{m}_{K^*}} (C_{10} - C_{10}') A_2(\hat{s}), \quad (27)$$

$$H(\hat{s}) = \frac{2\hat{m}_{K^*}}{\hat{s}} (C_{10} - C_{10}') (A_3(\hat{s}) - A_0(\hat{s})) + \frac{\hat{m}_{K^*}}{\hat{m}_l(\hat{m}_b + \hat{m}_s)} (C_{Q_2} - C_{Q_2}') A_0(\hat{s}) \quad (28)$$

$$S_2(\hat{s}) = \frac{1}{(\hat{m}_b + \hat{m}_s)} A_0(\hat{s}) (C_{Q_1}' - C_{Q_1}), \quad (29)$$

where $f_-(s) = \frac{m_B^2 - m_K^2}{s} (f_0(s) - f_+(s))$, $A_3(s) = \frac{m_B + m_{K^*}}{2m_{K^*}} A_1(s) - \frac{m_B - m_{K^*}}{2m_{K^*}} A_2(s)$. The above results reduce to those in ref. [13] if all $C_i' = 0$, as expected. It is worth to note that the final term in eq. (15) vanishes if one does not include the NHB contributions.

A. The dilepton invariant mass spectra and differential FBA

The kinematic variables $\hat{s}, \hat{u}$ are defined as:

$$\begin{aligned} \hat{s} &= \hat{q}^2 = (\hat{p}_+ + \hat{p}_-)^2, \\ \hat{u} &= (\hat{p}_B - \hat{p}_-)^2 - (\hat{p}_B - \hat{p}_+)^2 \end{aligned} \quad (30)$$

Here we choose the center of mass frame of the dileptons as the frame of reference, in which the leptons move back to back, and the momentum of B meson makes an angle θ with that of l^+ . $\hat{u}$ can be written in terms of θ :

$$\begin{aligned}\hat{u} &= -\hat{u}(\hat{s}) \cdot \cos \theta \equiv -\hat{u}(\hat{s})z, z = \cos \theta, \\ \hat{u}(\hat{s}) &= \sqrt{\lambda(1 - 4\frac{\hat{m}_l^2}{\hat{s}})}, \mathcal{D} = \sqrt{1 - 4\frac{\hat{m}_l^2}{\hat{s}}}, \\ \lambda &= 1 + \hat{m}_{K^{(*)}}^4 + \hat{s}^2 - 2\hat{s} - 2\hat{m}_{K^{(*)}}^2(1 + \hat{s})\end{aligned}\quad (31)$$

The phase space is defined in terms of $\hat{s}$ and z :

$$(2\hat{m}_l)^2 \leq \hat{s} \leq (1 - \hat{m}_{K^{(*)}})^2, -1 \leq z \leq 1 \quad (32)$$

Keeping the lepton mass and integrating over $\hat{u}$ in the kinematic region, we can get the dilepton invariant mass spectra (IMS):

$$\frac{d\Gamma^{K(K^*)}}{d\hat{s}} = \frac{G_F^2 \alpha^2 m_B^5}{2^{10} \pi^5} |V_{tb} V_{ts}^*|^2 \hat{u}(\hat{s}) D^{K(K^*)}, \quad (33)$$

$$\begin{aligned}D^K &= (|A'|^2 + |C'|^2)(\lambda - \frac{\hat{u}(\hat{s})^2}{3}) + |S_1|^2(\hat{s} - 4\hat{m}_l^2) \\ &\quad + |C'|^2 4\hat{m}_l^2(2 + 2\hat{m}_K^2 - \hat{s}) + \text{Re}(C' D'^\dagger) 8\hat{m}_l^2(1 - \hat{m}_K^2) + |D'|^2 4\hat{m}_l^2 \hat{s},\end{aligned}\quad (34)$$

$$\begin{aligned}D^{K^*} &= \frac{|A|^2}{3} \hat{s} \lambda (1 + 2\frac{\hat{m}_l^2}{\hat{s}}) + \frac{|E|^2}{3} \hat{s} \hat{u}(\hat{s})^2 + |S_2|^2(\hat{s} - 4\hat{m}_l^2) \lambda \\ &\quad + \frac{1}{4\hat{m}_{K^*}^2} \left[|B|^2(\lambda - \frac{\hat{u}(\hat{s})^2}{3}) + 8\hat{m}_{K^*}^2(\hat{s} + 2\hat{m}_l^2) + |F|^2(\lambda - \frac{\hat{u}(\hat{s})^2}{3}) + 8\hat{m}_{K^*}^2(\hat{s} - 4\hat{m}_l^2) \right] \\ &\quad + \frac{\lambda}{4\hat{m}_{K^*}^2} \left[|C|^2(\lambda - \frac{\hat{u}(\hat{s})^2}{3}) + |G|^2 \left(\lambda - \frac{\hat{u}(\hat{s})^2}{3} + 4\hat{m}_l^2(2 + 2\hat{m}_{K^*}^2 - \hat{s}) \right) \right] \\ &\quad - \frac{1}{2\hat{m}_{K^*}^2} \left[\text{Re}(BC^\dagger)(1 - \hat{m}_{K^*}^2 - \hat{s})(\lambda - \frac{\hat{u}(\hat{s})^2}{3}) \right. \\ &\quad \left. + \text{Re}(FG^\dagger) \left((1 - \hat{m}_{K^*}^2 - \hat{s})(\lambda - \frac{\hat{u}(\hat{s})^2}{3}) + 4\hat{m}_l^2 \lambda \right) \right] \\ &\quad - 2\frac{\hat{m}_l^2}{\hat{m}_{K^*}^2} \lambda [\text{Re}(FH^\dagger) - \text{Re}(GH^\dagger)(1 - \hat{m}_{K^*}^2)] + |H|^2 \frac{\hat{m}_l^2}{\hat{m}_{K^*}^2} \hat{s} \lambda\end{aligned}\quad (35)$$

The differential FBA is defined as:

$$A_{FB}(\hat{s}) = \frac{-\int_0^{\hat{u}(\hat{s})} dz \frac{d^2\Gamma}{d\hat{s}d\hat{u}} + \int_{-\hat{u}(\hat{s})}^0 d\hat{u} \frac{d^2\Gamma}{d\hat{s}d\hat{u}}}{\int_0^{\hat{u}(\hat{s})} dz \frac{d^2\Gamma}{d\hat{s}d\hat{u}} + \int_{-\hat{u}(\hat{s})}^0 d\hat{u} \frac{d^2\Gamma}{d\hat{s}d\hat{u}}} \quad (36)$$

According to the definition, it is straightforward to obtain the expressions of FBA in the exclusive decays:

- $B \rightarrow Kl^+l^-$

$$\frac{dA_{FB}^K}{d\hat{s}} D^K = -2\hat{m}_l \hat{u}(\hat{s}) \text{Re}(S_1 A'^\dagger) \quad (37)$$

- $B \rightarrow K^*l^+l^-$

$$\frac{dA_{FB}^{K^*}}{d\hat{s}} D^{K^*} = \hat{u}(\hat{s}) \left\{ \hat{s} [\text{Re}(BE^\dagger) + \text{Re}(AF^\dagger)] + \frac{\hat{m}_l}{\hat{m}_{K^*}} [\text{Re}(S_2 B^\dagger)(1 - \hat{s} - \hat{m}_{K^*}^2) - \text{Re}(S_2 C^\dagger) \lambda] \right\} \quad (38)$$

Seen from the Eqs. (33), (34), (35), (37) and (38), the functions $D'(\hat{s})$, $S_1(\hat{s})$, $H(\hat{s})$, and $S_2(\hat{s})$, which come from the contribution of NHBs, enter the IMS and FBA. Hence, the effects of NHBs will manifest themselves in the numerical results of these formula. In particular, Eq. (37) shows that the FBA in $B \rightarrow Kl^+l^-$ vanishes if there is no NHB contributions and from Eq. (38) it follows that the NHB contributions change the position of the zero-point of the FBA in $B \rightarrow K^*l^+l^-$. As pointed out in ref. [20], in an untagged sample, the FB asymmetry for unpolarized leptons vanishes. Once the flavor of the decaying b-quark is tagged, one can measure the unpolarized FB asymmetry which is an important observable to discriminate new physics from the SM, as we noted above.

B. The lepton polarization

In this subsection, we will present the analytical expressions of lepton polarization. We define the three orthogonal unit vectors in the center mass frame of dilepton as

$$\begin{aligned}\hat{e}_L &= \vec{p}_+, \\ \hat{e}_N &= \frac{\vec{p}_K \times \vec{p}_+}{|\vec{p}_K \times \vec{p}_+|}, \\ \hat{e}_T &= \hat{e}_N \times \hat{e}_L,\end{aligned}\tag{39}$$

which are related to the spin of lepton by a Lorentz boost. Then, the decay width of the $B \rightarrow K^{(*)}l^+l^-$ decay for any spin direction $\hat{n}$ of the lepton, where $\hat{n}$ is a unit vector in the dilepton center mass frame, can be written as:

$$\frac{d\Gamma(\hat{n})}{d\hat{s}} = \frac{1}{2} \left(\frac{d\Gamma}{d\hat{s}} \right)_0 [1 + (P_L \hat{e}_L + P_N \hat{e}_N + P_T \hat{e}_T) \cdot \hat{n}]\tag{40}$$

where the subscript "0" denotes the unpolarized decay width, P_L and P_T are the longitudinal and transverse polarization asymmetries in the decay plane respectively, and P_N is the normal polarization asymmetry in the direction perpendicular to the decay plane.

The lepton polarization asymmetry P_i can be obtained by calculating

$$P_i(\hat{s}) = \frac{d\Gamma(\hat{n} = \hat{e}_i)/d\hat{s} - d\Gamma(\hat{n} = -\hat{e}_i)/d\hat{s}}{d\Gamma(\hat{n} = \hat{e}_i)/d\hat{s} + d\Gamma(\hat{n} = -\hat{e}_i)/d\hat{s}}.\tag{41}$$

By a straightforward calculation, we get

- $B \rightarrow Kl^+l^-$

$$P_L^K D^K = \frac{4}{3} \mathcal{D} \{ \lambda \text{Re}(A' C'^\dagger) - 3\hat{m}_l(1 - \hat{m}_K^2) \text{Re}(C'^\dagger S_1) - 3\hat{m}_l \hat{s} \text{Re}(D'^\dagger S_1) \},\tag{42}$$

$$P_N^K D^K = \frac{\pi \sqrt{\hat{s}} \hat{u}(\hat{s})}{2} \left\{ -\text{Im}(A' S_1^\dagger) + 2\hat{m}_l \text{Im}(C' D'^\dagger) \right\},\tag{43}$$

$$P_T^K D^K = -\frac{\pi \sqrt{\lambda}}{\sqrt{\hat{s}}} \left\{ \hat{m}_l [(1 - \hat{m}_K^2) \text{Re}(A' C'^\dagger) + \hat{s} \text{Re}(A' D'^\dagger)] + \frac{(\hat{s} - 4\hat{m}_l^2)}{2} \text{Re}(C' S_1^\dagger) \right\}\tag{44}$$

- $B \rightarrow K^* l^+ l^-$

$$\begin{aligned}P_L^{K^*} D^{K^*} &= \mathcal{D} \left\{ \frac{2\hat{s}\lambda}{3} \text{Re}(A E^\dagger) + \frac{(\lambda + 12\hat{m}_{K^*}^2 \hat{s})}{3\hat{m}_{K^*}^2} \text{Re}(B F^\dagger) \right. \\ &\quad - \frac{\lambda(1 - \hat{m}_{K^*}^2 - \hat{s})}{3\hat{m}_{K^*}^2} \text{Re}(B G^\dagger + C F^\dagger) + \frac{\lambda^2}{3\hat{m}_{K^*}} \text{Re}(C G^\dagger) \\ &\quad \left. + \frac{2\hat{m}_l \lambda}{\hat{m}_{K^*}} [\text{Re}(F S_2^\dagger) - \hat{s} \text{Re}(H S_2^\dagger) - (1 - \hat{m}_{K^*}^2) \text{Re}(G S_2^\dagger)] \right\},\end{aligned}\tag{45}$$

$$\begin{aligned}P_N^{K^*} D^{K^*} &= \frac{-\pi \sqrt{\hat{s}} \hat{u}(\hat{s})}{4\hat{m}_{K^*}} \left\{ \frac{\hat{m}_l}{\hat{m}_{K^*}} [\text{Im}(F G^\dagger)(1 + 3\hat{m}_{K^*}^2 - \hat{s}) \right. \\ &\quad + \text{Im}(F H^\dagger)(1 - \hat{m}_{K^*}^2 - \hat{s}) - \text{Im}(G H^\dagger) \lambda] \\ &\quad + 2\hat{m}_{K^*} \hat{m}_l [\text{Im}(B E^\dagger) + \text{Im}(A F^\dagger)] \\ &\quad \left. - (1 - \hat{m}_{K^*}^2 - \hat{s}) \text{Im}(B S_2^\dagger) + \lambda \text{Im}(C S_2^\dagger) \right\},\end{aligned}\tag{46}$$

$$\begin{aligned}P_T^{K^*} D^{K^*} &= \frac{\pi \sqrt{\lambda} \hat{m}_l}{4\sqrt{\hat{s}}} \left\{ 4\hat{s} \text{Re}(A B^\dagger) \right. \\ &\quad + \frac{(1 - \hat{m}_{K^*}^2 - \hat{s})}{\hat{m}_{K^*}^2} [-\text{Re}(B F^\dagger) + (1 - \hat{m}_{K^*}^2) \text{Re}(B G^\dagger) + \hat{s} \text{Re}(B H^\dagger)] \\ &\quad + \frac{\lambda}{\hat{m}_{K^*}^2} [\text{Re}(C F^\dagger) - (1 - \hat{m}_{K^*}^2) \text{Re}(C G^\dagger) - \hat{s} \text{Re}(C H^\dagger)] \\ &\quad \left. + \frac{(\hat{s} - 4\hat{m}_l^2)}{\hat{m}_{K^*} \hat{m}_l} [(1 - \hat{m}_{K^*}^2 - \hat{s}) \text{Re}(F S_2^\dagger) - \lambda \text{Re}(G S_2^\dagger)] \right\}\end{aligned}\tag{47}$$

One can see from Eq. (43) that $P_N = 0$ for the decay $B \rightarrow Kl^+l^-$ in the SM because $C'_{10} = 0$ in the approximation of $m_s/m_b = 0$, $C'_{Q_{1,2}} = 0$, and C_{10} is real in the SM. Thus, a non zero normal polarization asymmetry in $B \rightarrow Kl^+l^-$ would signal the existence of new physics.




FIG. 1: The mass spectrum versus m_0 for fixed $M_{1/2}=200$ GeV, $\tan\beta=40$, $\delta_{23RR}^d=(0.04-0.03i)$, and $\text{sign}(\mu)=+1$ without the constraints from the low energy experiments imposed. (a) is for $A_0 = 0$. (b) is for $A_0=-1000$.

IV. MASS SPECTRA AND THE PERMITTED PARAMETER SPACE

To see the impact of the induced off-diagonal elements in the mass matrix of the right-handed down-type squarks on B rare decays and simplify the analysis, we assume that at the GUT scale (M_G) all sfermion mass matrices except the right-handed down-type squark mass matrix are flavor diagonal and all diagonal elements are approximately universal and equal to m_0^2 . The 2-3 matrix element of right-handed down-type squark mass matrix is parameterized by $\delta_{23}^{dRR} \equiv \frac{(M_{dRR}^2)_{23}}{m_0^2}$ which can be treated as a free parameter of order one. Furthermore, we have a universal gaugino mass $M_{1/2}$, a universal trilinear coupling A_0 and a universal bilinear coupling B_0 at M_G . Taking account to the radiative electro-weak (EW) symmetry breaking, finally we have five parameters ($m_0, M_{1/2}, A_0, \delta_{23}^{dRR}, \tan\beta$) plus the sign of μ as the initial conditions for solving the renormalization group equations (RGEs).

We require the lightest neutralino to be the lightest supersymmetric particle (LSP) and use several experimental limits to constraint the parameter space, including 1) the width of the decay $Z \rightarrow \chi_1^0 \chi_1^0$ is less than 4.3 MeV, and branching ratios of $Z \rightarrow \chi_1^0 \chi_2^0$ and $Z \rightarrow \chi_2^0 \chi_2^0$ are less than 1×10^{-5} , where χ_1^0 is the lightest neutralino and χ_2^0 is the other neutralino; 2) the mass of light neutral Higgs can not be lower than 111 GeV as the present experiments required; 3) the mass of lighter chargino must be larger than 94 GeV as given by the Particle Data Group [40]; 4) sneutrinos are larger than 94 GeV; 5) seletrons are larger than 73 GeV; 6) smuons larger than 94 GeV; 7) staus larger than 81.9 GeV.

In the numerical calculation, we use the revised ISAJET. We find that the parameter δ_{23}^{dRR} does not receive any significant correction and the diagonal entries of mass matrices are significantly corrected, which is in agreement with the results in Ref. [41]. We scan $m_0, M_{1/2}$ in the range (100, 800) GeV for given values of $A_0, \tan\beta$ and $\text{sign}(\mu)=+1^*$, with the constraints from the relevant low energy experiments such as $B \rightarrow X_s \gamma, B_s \rightarrow \mu^+ \mu^-$, etc. (for the detailed discussions of constraints, see VB).

For an illustration, we present the mass spectra without and with the constraints from the low energy experiments in Figs. 1 and 2, respectively, where (a) and (b) are for $A_0 = 0, -1000$ GeV respectively. One can see from the Figs. 1, 2 that the mass spectrum lifts when A_0 increases and when the constraints from the low energy experiments are imposed the masses of particles are larger than those without the constraints, as expected.

V. NUMERICAL ANALYSIS

In this section, we will discuss the numerical results and make an analysis.

* In the case of $\text{sign}(\mu)=-1$, the constraint from $B \rightarrow X_s \gamma$ on the parameter space is too stringent, in particular, for large $\tan\beta$ [32, 44, 45]

TABLE I: The Wilson coefficients for the two cases in the $SUSY SO(10)$. The SM values also are listed for comparison. The values in bracket are for $l = \tau$.

A_0	C_{Q_1}	C'_{Q_1}	C_{Q_2}	C'_{Q_2}
SM	0	0	0	0
$A_0 = 0$	0.074+0.I(1.252+0.001I)	-0.013+0.008I(-0.213+0.128I)	-0.075+0.I(-1.267-0.001I)	-0.013+0.008I(-0.216+0.129I)
$A_0 = -1000$	0.106+0.I(1.775+0.002I)	-0.247+0.242I(-4.148+4.074I)	-0.107+0.I(-1.797-0.002I)	-0.250+0.246I(-4.202+4.128I)

A_0	C_7^{eff}	$C_7'^{eff}$	C_9	C_9'	C_{10}	C_{10}'
SM	-0.313	0	+4.344	0	-4.669	0
$A_0 = 0$	-0.225-0.I	-0.02-0.01I	4.277+0.I	-0.-0.002I	-4.717-0.I	-0.001+0.019I
$A_0 = -1000$	-0.219+0.I	0.039-0.038I	4.275+0.I	0.011+0.072I	-4.732-0.I	-0.075-0.670I

and consequently the parameter space of the model. At the same time we require that predicted branching ratios of $B \rightarrow X_s \mu^+ \mu^-$ and $B \rightarrow K^{(*)} \mu^+ \mu^-$ falls within 1 σ experimental bounds.

We also impose the current experimental lower bound $\Delta M_s > 14.4 ps^{-1}$ [38]. The $\delta_{23}^{dLR(RL)}$ contribution to ΔM_s is small because it is constrained to be order of 10^{-2} by $Br(B \rightarrow X_s \gamma)$. The dominant contribution to ΔM_s comes from $\delta_{23}^{dLL(RR)}$ insertion with both constructive and destructive effects compared with the SM contribution, where the too large destructive effect is ruled out, because SM prediction is only slightly above the present experiment lower bound.

Furthermore, as analyzed in ref. [41], there is the correlation between flavor changing squark and slepton mass insertions in SUSY GUTs. The correlation leads to a bound on δ_{23}^{dRR} from the rare decay $\tau \rightarrow \mu \gamma$. We update the analysis with latest BELLE upper bound of $Br(\tau \rightarrow \mu \gamma) < 3.1 \times 10^{-7}$ [42] at 90% confidence level in the SUSY SO(10) model.

C. The Numerical results and Conclusions

We will focus on the parameter space at large $\tan\beta$. The reason for this is that in the large $\tan\beta$ region of parameter space the contributions of NHB exchange become very important for quark level semi-leptonic transitions $b \rightarrow sl^+ l^-$ when the final state lepton is either a muon or tau [32, 45]. In numerical calculations, we take $\tan\beta=40$, $\text{sign}(\mu)=+1$, and $A_0 = 0, -1000$ and get the sparticle mass spectrum and mixing at the EW scale. Using the resulted Wilson coefficients, we calculate the IMS, FBA and polarization asymmetries of processes $B \rightarrow K^{(*)} l^+ l^-$ under the constraints on δ_{23}^{dRR} from the all relevant experiments as discussed in subsection VB, whose phase varies from 0 to 2π .

The Wilson coefficients $C_{Q_1}^{(\prime)}$ and $C_{Q_2}^{(\prime)}$ come from NHB exchanging. Specially, we are interested in the case of maximal enhancements of $C_{Q_{1,2}}^{(\prime)}$. Through scanning the parameter space under constraints, we find, for $A_0 = 0$, when $m_0=800\text{GeV}$, $M_{1/2}=400\text{GeV}$ (for $A_0=-1000$, $M_{1/2}=500\text{GeV}$), $C_{Q_{1,2}}^{(\prime)}$ have maximal values. The obtained $C_{Q_{1,2}}^{(\prime)}$ and other relevant Wilson coefficients in the two cases and in SM are listed in Table I. From Table I, we can know: (i) the NHB contributions in the case of $A_0 = -1000$ are larger than those in the case of $A_0 = 0$ except for C_9 ; (ii) the Wilson coefficients $C_{Q_{1,2}}^{(\prime)}$ of primed operators are dominant, which is due to the presence of δ_{23}^{dRR} of order one at the high scale in the SUSY SO(10) model, and their imaginary parts are sizable, which contribute to the normal polarization, a T violating observable.

The figures for the dependence of observable on $\hat{s}$ with and without long-distance contributions are presented in **Fig.(3-7)** in the case of $A_0 = -1000$, where the solid lines denote the all contributions ($W^\pm, H^\pm$, chargino, gluino, neutrino propagated in the loop) including the NHB contributions; the dot lines present the SM contribution plus only the NHB contributions, and the dot-dashed lines are for the SM contribution. We also calculated the dependence of observable on $\hat{s}$ in the case of $A_0 = 0$ and give the results for $A_0 = 0$ when the two cases have a sizable difference.

The IMS of the process $B \rightarrow K^{(*)} l^+ l^-$ is given in **Fig.3**, where the left two figures are for $B \rightarrow K^{(*)} \mu^+ \mu^-$ and the right for $B \rightarrow K^{(*)} \tau^+ \tau^-$. For the case of $B \rightarrow K \mu^+ \mu^-$, we can see that, at the low $\hat{s}$ region, there are some enhancement from NHB contributions. Compared to the decay $B \rightarrow K \mu^+ \mu^-$, the IMS of $B \rightarrow K^* \mu^+ \mu^-$ deviates from the SM prediction sizably in the whole region of $\hat{s}$. Nevertheless, the SUSY effects are small compared with the SM for the decay $B \rightarrow K^{(*)} \tau^+ \tau^-$.

The **Fig. 4** is for the FBA ($dA/d\hat{s}$) of the decay $B \rightarrow K^{(*)} l^+ l^-$, where the left two figures are for $l = \mu$ and the right for $l = \tau$, like that in Fig.3. As it is known, the FBA ($dA/d\hat{s}$) of $B \rightarrow Kl^+ l^-$ in the SM is zero. Since FBA arises in the SUSY models only when NHB effects are taken into account, it provides a good probe to test these effects. Our numerical results show that the average of FBA in $B \rightarrow K \mu^+ \mu^-$ can reach only 0.001 which is too small to be observed. The average of FBA in $B \rightarrow K \tau^+ \tau^-$ can reach -0.1 and 0.05 for the case of $A_0 = 0$ and $A_0 = -1000$, respectively. (The reason why FBA in the two cases has the opposite sign is that the sign of function S_1 in these two cases is opposite.) So, the $10^{10} - 10^{11}$ $B_d \bar{B}_d$ pairs per year, which is in the designed range in the future super B factories with B hadrons $10^{10} - 10^{12}$ per year [43], is needed in order to observe the FBA with good accuracy. Our

results show that the SUSY effects show up at the low $\hat{s}$ region for the FBA of $B \rightarrow K^* \mu^+ \mu^-$ and the deviation from SM is 0.05 or so. It is worth to note that there is a sizable change of the position of the zero-point of the FBA in $B \rightarrow K^* \mu^+ \mu^-$ in the SUSY SO(10) model, as it can be seen in Fig.5, which could be tested in the future experiments with high precision. For FBA in $B \rightarrow K^* \tau^+ \tau^-$, the deviation from the SM is about several percent. The average of FBA of $B \rightarrow K^* \tau^+ \tau^-$ can reach 0.3 in the case of $A_0 = 0$. To observe the FBA in $B \rightarrow K^* \tau^+ \tau^-$ decay at 1σ level, the required number of events is 1.1×10^8 . The number of $B\bar{B}$ pairs that is expected to be produced at B factories is about $N \simeq 5 \times 10^8$. Therefore the FBA in $B \rightarrow K^* \tau^+ \tau^-$ could be observed at B factories. Hence, with the enhancement of experimental precision and statistics, the measurements of FBA would provide more data and effectively pin to the NP effects.

Now, we turn to discuss the lepton polarization. We present the longitudinal, transverse and normal polarization in the Fig.(5,6,7) for $B \rightarrow K^{(*)} l^+ l^-$ decay. As it can be seen in Fig. 5 and 6, the $dL/d\hat{s}$ of $B \rightarrow K^{(*)} \mu^+ \mu^-$ is not sensitive to the NHB effects, while for $dT/d\hat{s}$ of $B \rightarrow K(K^*) \mu^+ \mu^-$, the deviation from SM can reach 0.1 (0.05). As it is expected, the contribution from the $\tau^+ \tau^-$ channel is much larger than that from the $\mu^+ \mu^-$ one. For $B \rightarrow K^{(*)} \tau^+ \tau^-$, the NHB contributions are manifest and dominant, and both $dL/d\hat{s}$ and $dT/d\hat{s}$ are significantly different from SM. And the $dL/d\hat{s}$ of $B \rightarrow K \tau^+ \tau^-$ can even reach 0.6. Thus, the NHB effects are sensitive to $B \rightarrow K^{(*)} \tau^+ \tau^-$ and will be observable at B factories.

The $dN/d\hat{s}$ of $B \rightarrow K^{(*)} l^+ l^-$ decay are given in Fig. 7. The average of $dN/d\hat{s}$ can reach several percent for $B \rightarrow K \mu^+ \mu^-$ which could be observed in the future super B factories, while it is the order of 10^{-3} for $B \rightarrow K^* \mu^+ \mu^-$ which can not be observed even in designed super B factories. The average of $dN/d\hat{s}$ in $B \rightarrow K \tau^+ \tau^-$ is 0.05 or so. For $B \rightarrow K^* \tau^+ \tau^-$, the deviation from SM is a few percent. As noted above, the Wilson coefficient C_{10} is real and $C'_{10}, C_{Q_i}^{(\prime)} = 0$ (precisely speaking, they are negligibly small) in the SM so that $dN/d\hat{s} = 0$ in $B \rightarrow K \mu^+ \mu^-$ in the SM. It is still true in the minimal super gravity model (mSUGRA) and SUSY models with real universal boundary conditions at the high scale [13]. In the SUSY SO(10) model we considered, the complex flavor non-diagonal down-type squark mass matrix element of 2nd and 3rd generations of order one at the GUT scale induces the complex couplings which lead to the complex Wilson coefficients and consequently the non zero normal polarization of $B \rightarrow K \mu^+ \mu^-$. Therefore, the measurements of the CP violating (as usual, the CPT invariance is assumed in the paper) normal polarization in $B \rightarrow K l^+ l^-$ could discriminate the SUSY SO(10) model (and other SUSY models with the flavor non diagonal complex couplings) from the SM and mSUGRA.

In summary, we have carried out a study of SUSY effects, in particular, the neutral Higgs bosons contributions to the IMS, FBA and polarization, in the exclusive decay $B \rightarrow K^{(*)} l^+ l^-$ ($l = \mu, \tau$) in the SUSY SO(10) model, taking account of the constraints from existing experimental data such as $b \rightarrow s\gamma$, ΔM_s , $Br(B \rightarrow K^{(*)} \mu^+ \mu^-)$, $\tau \rightarrow \mu\gamma$ as well as the upper bound of $Br(B_s \rightarrow \mu^+ \mu^-)$. Our main findings can be summarized as follows:

- The IMS of the process $B \rightarrow K^{(*)} \mu^+ \mu^-$ can sizably deviate from the SM.
- The FBA comes only from NHB contributions in $B \rightarrow K l^+ l^-$ and its average for $l = \mu$ is nonzero but too small to be observed. However for $B \rightarrow K \tau^+ \tau^-$, it is the order of 10%, which should be within the luminosity reach of coming B factories. The SUSY effects show up at the low $\hat{s}$ region for the FBA of $B \rightarrow K^* \mu^+ \mu^-$ and the deviation from SM is 0.05 or so. Moreover, there is a sizable change of the position of the zero-point of the FBA in $B \rightarrow K^* \mu^+ \mu^-$, which can be used to discriminate the model from the SM.
- The average of $dN/d\hat{s}$ can reach several percent for $B \rightarrow K \mu^+ \mu^-$ and it is 0.05 or so for $B \rightarrow K \tau^+ \tau^-$, which could be measured in the future super B factories and provide a useful information to probe new physics and discriminate different models.
- The longitudinal polarization, $dL/d\hat{s}$, of $B \rightarrow K^{(*)} \mu^+ \mu^-$ is not sensitive to the NHB effects. However, for the transverse polarization, $dT/d\hat{s}$, of $B \rightarrow K(K^*) \mu^+ \mu^-$, the deviation from SM can reach 0.1 (0.05) which could be seen in B factories. For $B \rightarrow K^{(*)} \tau^+ \tau^-$, the NHB contributions are manifest and dominant, and both $dL/d\hat{s}$ and $dT/d\hat{s}$ are significantly different from SM. And the $dL/d\hat{s}$ of $B \rightarrow K \tau^+ \tau^-$ can even reach 0.6, which can be measured in B factories.

Therefore, the experimental investigation of observable, in particular, FBA and the polarization components, in the $B \rightarrow K^{(*)} l^+ l^-$ decays in the present B factories and future super B factories can be used to search for SUSY effects, in particular, NHB effects, in SUSY grand unification models.

Acknowledgement

One of the authors (W.-J. Li) would like to thank Dr. X.-H. Wu for discussions during the work. The work was supported in part by the National Nature Science Foundation of China.

References

-
- [1] A.Ishikawa, (Belle Collaboration), Lecture given at ICHEP 2004.
- [2] A. Ishikawa *et al.*, BELLE Collaboration, Phys. Rev. Lett. **91**, 261601 (2003); B. Aubert *et al.*, (BABAR Collaboration), Phys. Rev. Lett. **91**, 221802 (2003).
- [3] P. Ball and R. Zwicky, hep-ph/0406232; M.A. Ivanov and V.E. Lyubovitskij, Lectures given at International School on Heavy Quark Physics, Dubna, Russia, 27 May - 5 Jun 2002; W. Jaus and D. Wyler, Phys. Rev. **D 41** (1990) 3405; P. Colangelo *et al.*, Phys. Lett. **B317** (1993) 183; Ceng *et al.*, Phys. Rev. **D 54** (1996) 3656; Amand Faessler *et al.*, EPJ C, **4**, C18, 1C33 (2002).
- [4] P. Colangelo *et al.*, Phys. Rev. **D 53** (1996) 3672 [arXiv:hep-ph/9510403]; Erratum, *ibid*, **D 57** (1998) 3186.
- [5] I.I. Balitsky, V.M. Braun and A.V. Kolesnichenko, Nucl. Phys. **B312** (1989) 509; V.L. Chernyak and I.R. Zhitnitsky, Nucl. Phys. **B345** (1990) 137; V.M. Braun, Preprint NORDITAC98C1CP [arXiv:hep-ph/9801222]; A. Khodjamirian and R. Ruckl, [arXiv:hep-ph/9801443]; P. Ball and V.M. Braun, Phys. Rev. **D 58** (1998) 094016 [arXiv:hep-ph/9805422]; Nucl. Phys. **B543** (1999) 201 [arXiv:hep-ph/9810475]; P. Ball *et al.*, Nucl. Phys. **B529** (1998) 323 [arXiv:hep-ph/9802299]; P. Ball, JHEP **09** (1998) 005 [arXiv:hep-ph/9802394]; JHEP **01** (1999) 010 [arXiv:hep-ph/9812375].
- [6] C.Q. Geng and C.P. Kao, Phys. Rev. **D 54** (1996) 5636 [arXiv:hep-ph/9608466]; Phys. Rev. **D 57** (1998) 4479.
- [7] S. Fukaea, C.S. Kimb, T. Morozumi, and T. Yoshikawac, Phys. Rev. **D 59** (1999) 074013 [arXiv:hep-ph/9807254]; T.M. Aliev, C. S. Kim and Y. G. Kim, Phys. Rev. **D 62**, 014026 (2000) [arXiv:hep-ph/9910501]; T. M. Aliev, *et al.*, Nucl. Phys. **B607** (2001) 305-325 [arXiv:hep-ph/0009133]; T.M. Aliev, M.K. Cakmak, A. Özpineci, M. Savci, Phys. Rev. **D 64** (2001) 055007 [arXiv:hep-ph/0103039].
- [8] Benjamin Grinstein, Dan Pirjol, CTP-MIT-3490, [arXiv:hep-ph/0404250].
- [9] T.M. Aliev, M. Savci, Phys. Lett. **B481** (2000) 275-286 [arXiv:hep-ph/0003188].
- [10] L.T. Handoko, C.S. Kim, T. Yoshikawa, Phys. Rev. **D 65** (2002) 077506 [arXiv:hep-ph/0112149].
- [11] A. Ali, E. Lunghi, C. Greub, G. Hiller, DESY 01-217, BUTP-01-21, SLAC-PUB-9076 [arXiv:hep-ph/0112300].
- [12] T.M. Aliev, M. Savci, A. Ozpineci, H. Koru, J.Phys. G24 (1998) 49-65 [arXiv:hep-ph/9705222].
- [13] Qi-Shu Yan, Chao-Shang Huang, Liao Wei, Shou-Hua Zhu, Phys. Rev. **D 62** (2000) 094023 [arXiv:hep-ph/0004262].
- [14] A. Ali, P. Ball, L.T. Handoko, G. Hiller, Phys. Rev. **D 61** (2000) 074024 [arXiv:hep-ph/9910221].
- [15] F. Krüger, E. Lunghi, Phys. Rev. **D 63** (2001) 014013 [arXiv:hep-ph/0008210].
- [16] C.S. Huang, Nucl.Phys. Proc. Suppl. **93** (2001) 73-78 [arXiv:hep-ph/0009149]; C. Bobeth, T. Ewerth, F. Kruger, J. Urban, Phys. Rev. **D 64** (2001) 074014 [arXiv:hep-ph/0104284].
- [17] Guray Erkol, Gursevil Turan, Nucl. Phys. **B635** (2002) 286-308 [arXiv:hep-ph/0204219].
- [18] S. R. Choudhury *et al.*, Phys. Rev. **D 69** (2004) 054018 [arXiv:hep-ph/0307276].
- [19] S. Rai Choudhury *et al.*, Phys. Rev. **D 68**, 054016 (2003) [arXiv:hep-ph/0304084].
- [20] Wafia Bensalam, David London, Nita Sinha, Rahul Sinha, Phys. Rev. **D 67** (2003) 034007 [arXiv:hep-ph/0209228].
- [21] T.M. Aliev, V. Bashiry, M. Savci, Eur. Phys. J. **C35** (2004) 197-206 [arXiv:hep-ph/0311294]; JHEP **0405** (2004) 037 [arXiv:hep-ph/0403282]; A.S. Cornell, Naveen Gaur, [arXiv:hep-ph/0408164].
- [22] V. Barger *et al.*, Phys. Rev. Lett. **82** (1999) 2640 [arXiv:astro-ph/9810121]; E. Ili *et al.*, Phys. Rev. Lett. **85** (2000) 1166 [arXiv:hep-ph/0002053]; John N. Bahcall *et al.*, Phys. Rev. Lett. **90**, 131301 (2003) [arXiv:astro-ph/0212331]; John N. Bahcall and M. H. Pinsonneault, Phys. Rev. Lett. **92**, 121301 (2004) [arXiv:astro-ph/0402114]; Sandhya Choubey and Probir Roy, Phys. Rev. Lett. **93**, 021803 (2004) [arXiv:hep-ph/0310316].
- [23] K.S. Babu and S.M. Barr, Phys. Lett. **B381** (1996) 202 [arXiv:hep-ph/9511446]; C.H. Albright, K.S. Babu, and S.M. Barr, Phys. Rev. Lett. **81** (1998) 1167 [arXiv:hep-ph/9802314]; J. Sato and T. Yanagida, Phys. Lett. **B430** (1998) 127 [arXiv:hep-ph/9710516]; N. Irges, S. Lavignac, and P. Ramond, Phys. Rev. **D 58** (1998) 035003 [arXiv:hep-ph/9802334].
- [24] Mu-Chun Chen, K.T. Mahanthappa, Phys. Rev. **D62** 113007 (2000), [arXiv:hep-ph/0005292].
- [25] D. Chang, A. Masiero and H. Murayama, Phys. Rev. **D67** (2003) 075013 [arXiv:hep-ph/0205111].
- [26] B. Bajc, G. Senjanović and F. Vissani, [arXiv:hep-ph/0210207]; H.S. Goh, R.N. Mohapatra and S.-P. Ng, [arXiv:hep-ph/0303055].
- [27] X.-J. Bi, Y.-B. Dai and X.-Y. Qi, Phys. Rev. **D63** 096008 (2001); X.-J. Bi and Y.-B. Dai, Phys. Rev. **D66** 076006 (2002).
- [28] Mu-Chun Chen, K.T. Mahanthappa, Phys. Rev. **D65** 053010 (2002), [arXiv:hep-ph/0106093]; Mu-Chun Chen, K.T. Mahanthappa, Phys. Rev. **D68** 017301 (2003), [arXiv:hep-ph/0212375]; Mu-Chun Chen and K.T. Mahanthappa, Int.J.Mod.Phys. **A18**:5819-5888, 2003 [arXiv:hep-ph/0305088]; Mu-Chun Chen, K.T. Mahanthappa, [arXiv:hep-ph/0409096].
- [29] Sebastian Jager, Ulrich Nierste, FERMILAB-Conf-03/394-T, [arXiv:hep-ph/0312145].
- [30] Y.-B. Dai, C.-S. Huang, W.-J. Li, X.-H. Wu, [arXiv:hep-ph/0407263].
- [31] T. Goto *et al.*, Phys. Rev. **D 55** (1997) 4273; T. Goto, Y. Okada and Y. Shimizu, Phys. Rev. **D 58** (1998) 094006; S. Bertolini, F. Borzynatu, A. Masiero and G. Ridolfi, Nucl. Phys. **B353** (91) 591.
- [32] Y.-B. Dai, C.-S. Huang, H.-W. Huang, Phys. Lett. **B390**:257-262, 1997; Erratum-*ibid*. **B513**:429-430, 2001 [arXiv:hep-ph/9607389].
- [33] Chao-Shang Huang, Xiao-Hong Wu, Nucl. Phys. **B657** (2003) 304-332 [arXiv:hep-ph/0212220]; Jian-Feng Cheng, Chao-Shang Huang, Xiao-Hong Wu, hep-ph/0404055, to appear in Nucl. Phys. B.
- [34] A.J. Buras, M. Muenz, Phys. Rev. **D 52** (1995) 186-195 [arXiv:hep-ph/9501281].
- [35] D. Acosta *et al.*, (CDF Collaboration), [arXiv:hep-ex/0403032].
- [36] C.-S. Huang, W. Liao, Q.-S. Yan, S.-H. Zhu, Phys. Rev. **D63** (2001) 114021; *ibid*. **64** (2001) 059902(E) [arXiv:hep-ph/0006250]; For review papers, see, e.g., C.-S. Huang, Nucl. Phys. Proc. Suppl. **115**, 89 (2003); A. Dedes,




FIG. 3: The IMS of the process $B \rightarrow K^{(*)} l^+ l^-$ for $A_0 = -1000$. The solid line(black), dot line(green), and dashed-dot line(blue) represent the all contributions included, the SM contributions plus only the NHB contributions, and the SM contributions, respectively. Both the total (SD+LD) and the pure SD contributions are shown in order to compare. In the fig. we write "s" in stead of " $\hat{s}$ " for simplicity.

- Mod. Phys. Lett. **A18**, 2627 (2003) [arXiv:hep-ph/0309233].
- [37] The D_0 collaboration, D_0 Conference Note 4514.
- [38] A. Stocchi, Nucl. Phys. Proc. Suppl. **117**(2003) 145 [arXiv:hep-ph/0211245].
- [39] A. Kagan, [arXiv:hep-ph/9806266]; T.E. Coan *et al.*, (CLEO Collaboration), Phys. Rev. Lett. **80**, 1150 (1998) [arXiv:hep-ex/9710028].
- [40] K. Hagiwara *et al.*, Phys. Rev. **D66**, 010001 (2002).
- [41] M. Ciuchini *et al.*, Phys. Rev. Lett. **92** (2004) 071801.
- [42] K.Abe *et al.*, Belle collaboration, Phys. Rev. Lett. **92** (2004) 171802 [arXiv:hep-ex/0310029].
- [43] T.E. Browder and A. Soni, hep-ph/0410192, and references therein.
- [44] J.L. Lopez, D.V. Nanopoulos, X. Wang and A. Zichichi, Phys. Rev. **D51**, 147 (1995) [arXiv:hep-ph/9406427].
- [45] C.-S. Huang and Q.-S. Yan, Phys. Lett. **B442**(1998) 209 [arXiv:hep-ph/9803366]; C.-S. Huang, W. Liao, and Q.-S. Yan, Phys. Rev. **D59** (1999) 011701 [arXiv:hep-ph/9803460].
- [46] A. Stocchi, Nucl. Phys. Proc. Suppl. **117** (2003) 145 [arXiv:hep-ph/0211245].




FIG. 4: The FBA of the process $B \rightarrow K^{(*)} l^+ l^-$ for $A_0 = -1000$. The line conventions are the same as those in Fig. 3.




FIG. 5: The dL/ds of the process $B \rightarrow K^{(*)} l^+ l^-$ for $A_0 = -1000$. The line conventions are the same as those in Fig. 3.




FIG. 6: The dT/ds of the process $B \rightarrow K^{(*)} l^+ l^-$ for $A_0 = -1000$. The line conventions are the same as those in Fig. 3.




FIG. 7: The $dN/d\hat{s}$ of the process $B \rightarrow K^{(*)}l^+l^-$ for $A_0 = -1000$. The line conventions are the same as those in Fig. 3.